



QP - 173

40
V Semester B.A./B.Sc. Examination, March/April 2022
(Semester Scheme)
(CBCS) (F + R) (2016 – 17 and Onwards)
MATHEMATICS (Paper – V)



Time : 3 Hours

Max. Marks : 70

Instruction : Answer *all* questions.

PART – A

1. Answer **any five** questions.

(5×2=10)

- In a ring $(R, +, \cdot)$, show that $a \cdot (-b) = (-a) \cdot b = -(a \cdot b)$ for all $a, b \in R$.
- Define subring of a ring. Give an example.
- Give an example of
 - Commutative ring without unity.
 - A non commutative ring without unity.
- If $\phi(x, y, z) = x^2y^2z^2$ and $\vec{F} = 2x\hat{i} + y\hat{j} + 3z\hat{k}$ find $\vec{F} \cdot \nabla\phi$.
- Find the unit normal vector to the surface $x^2 - y^2 + z = 3$ at the point $(1, 0, 2)$.
- Evaluate : $\Delta^3[(1-x)(1-2x)(1-3x)]$.
- Write Lagrange's interpolation formula.
- Evaluate : $\int_0^1 \frac{dx}{1+x}$ using Trapezoidal rule, given

x	0	$\frac{1}{6}$	$\frac{2}{6}$	$\frac{3}{6}$	$\frac{4}{6}$	$\frac{5}{6}$	1
y	1	0.8571	0.75	0.6667	0.6	0.5455	0.5

PART – B

Answer **two full** questions.

(2×10=20)

- Prove that every field is an integral domain.
Is the converse of the above theorem is true ? Justify with example.
- Show that set $R = \{0, 1, 2, 3, 4, 5\}$ is a commutative ring w.r.t \oplus_6 and \otimes_6 as two compositions.

OR

P.T.O.



3. a) Prove that a ring R without zero divisors if and only if the cancellation laws holds.
- b) Show that necessary and sufficient condition for a non-empty subset S of a ring R to be a subring of R are
- $a - b \in S \quad \forall a, b \in S$
 - $ab \in S \quad \forall a, b \in S$.
4. a) Prove that an ideal S of the ring $(\mathbb{Z}, +, \cdot)$ is maximal if and only if S is generated by some prime integer.
- b) Find all the principal ideals of the ring $R = \{0, 1, 2, 3, 4, 5\}$ w.r.t \oplus_6 and \otimes_6 .
- OR
5. a) If $f : R \rightarrow R'$ be a homomorphism of R into R' , then show that $\text{Ker } f$ is an ideal of R .
- b) State and prove fundamental theorem of homomorphism.

PART – C

Answer **two full** questions.**(2×10=20)**

6. a) Find the directional derivative of $\phi(x, y, z) = xyz - xy^2z^3$ at the point $(1, 2, -1)$ in the direction of $\hat{i} - \hat{j} - 3\hat{k}$.
- b) If $\vec{F} = \text{grad}(x^3 + y^3 + z^3 - 3xyz)$, find $\text{div } \vec{F}$ and $\text{curl } \vec{F}$.
- OR
7. a) Find the values of 'a' and 'b' so that the surface $5x^2 - 2yz - 9x = 0$ may cut the surface $ax^2 + by^3 = 4$ orthogonally at $(1, -1, 2)$.
- b) If ϕ is a scalar point function and \vec{F} is a vector point function, prove that $\text{div}(\phi\vec{F}) = \phi(\text{div}\vec{F}) + (\text{grad } \phi) \cdot \vec{F}$.
8. a) If $\vec{u} = x^2\hat{i} + y^2\hat{j} + z^2\hat{k}$ and $\vec{v} = yz\hat{i} + zx\hat{j} + xy\hat{k}$. Show that $\vec{u} \times \vec{v}$ is a solenoidal vector.
- b) Show that $\vec{F} = (6xy + z^3)\hat{i} + (3x^2 - z)\hat{j} + (3xz^2 - y)\hat{k}$ is irrotational. Find ϕ such that $\vec{F} = \nabla\phi$.
- OR
9. a) Prove that :
- $\text{div}(\text{curl } \vec{F}) = 0$.
 - $\text{curl}(\text{grad } \phi) = 0$.
- b) For any vector field \vec{f} and \vec{g} prove that $\text{div}(\vec{f} \times \vec{g}) = \vec{g} \cdot \text{curl } \vec{f} - \vec{f} \cdot \text{curl } \vec{g}$.



PART – D

Answer any two full questions.

(2×10=20)

10. a) Use the method of separation of symbols. Prove that

$$u_0 + \frac{u_1x}{1!} + \frac{u_2x^2}{2!} + \dots = e^x \left[u_0 + x \frac{\Delta u_0}{1!} + x^2 \frac{\Delta^2 u_0}{2!} + \dots \right].$$

b) Obtain the function whose first difference is $6x^2 + 10x + 11$.

OR

11. a) Find a cubic polynomial which takes the following data and hence evaluate $f(4)$.

x	0	1	2	3
f(x)	1	2	1	10

b) Find $f(1.4)$ from the following data using difference table.

x	1	2	3	4	5
f(x)	10	26	58	112	194

12. a) Use Newton divided difference formula and find $f(8)$ from the following data :

x	1	3	6	11
f(x)	4	32	224	1344

b) Evaluate $\int_0^6 \frac{1}{1+x^2} dx$ by using Simpson's $\frac{3}{8}$ th rule by taking $n = 6$.

OR

13. a) By using Lagrange's interpolation formula, find $f(6)$ from the following data :

x	3	7	9	10
f(x)	168	120	72	63

b) Evaluate $\int_0^{0.6} e^{-x^2} dx$ by taking 6 subintervals by using Simpson's $\frac{1}{3}$ rd rule.
