



VI Semester B.A./B.Sc. Examination, Sept./Oct. 2022
(Semester Scheme)
(CBCS) (F+R) (2016-17 and Onwards)
MATHEMATICS – VIII

Time : 3 Hours

Max. Marks : 70

Instruction : Answer all Parts.

PART – A



1. Answer any five questions.

(5×2=10)

- Evaluate $\lim_{z \rightarrow 1+2i} (z^2 + 1)$.
- Find the locus of z satisfying $|z - i| \leq 3$.
- Show that $v = 3x^2y - y^3$ is harmonic.
- State Liouville's theorem.
- Define bilinear transformation.
- Verify Cauchy-Riemann equations for $f(z) = \sin x \cosh y + i \cos x \sinh y$.
- Find the real root of the equation $x^3 - 4x + 9 = 0$ in one step by bisection method.
- Write Euler's modified formula.

PART – B

Answer four full questions.

(4×10=40)

- Show that $|z + i|^2 - |z - i|^2 = 2$ represents a real axis.
 - State and prove necessary conditions for a function $f(z) = u + iv$ to be analytic.

OR

- Evaluate $\lim_{z \rightarrow \frac{i}{2}} \left[\frac{(2z^3 - 3)(4z + i)}{(iz - 1)^2} \right]$.

- Show that $f(z) = \log z$ is analytic and find $f'(z)$.



4. a) Find the analytic function $f(z) = u + iv$ given its real part $u = \left(r + \frac{1}{r}\right) \cos\theta$.
 b) Show that $u = e^x \sin y + x^2 - y^2$ is harmonic and find its harmonic conjugate.

OR

5. a) If $f(z)$ is analytic, show that $\left(\frac{\delta^2}{\delta x^2} + \frac{\delta^2}{\delta y^2}\right) |f(z)|^2 = 4 |f'(z)|^2$.
 b) Find the orthogonal trajectory of the family of curves $x^2 - y^2 - x = c$.
6. a) If $f(z)$ is analytic within and on a closed curve C of a simply connected region and $z = z_0$ is an interior point of C , prove that $\frac{1}{2\pi i} \int_C \frac{f(z)}{(z - z_0)} dz = f(z_0)$.
 b) Evaluate $\int_C \frac{z}{(z^2 + 1)(z^2 - 9)} dz$, where C is the circle $|z| = 2$.

OR

7. a) State and prove fundamental theorem of algebra.
 b) If C is the circle with centre 'a' and radius 'r' then show that
 i) $\int_C \frac{1}{(z - a)} dz = 2\pi i$
 ii) $\oint_C (z - a)^n dz = 0$, if $n \neq -1$.
8. a) Discuss the transformation $w = \sin z$.
 b) Show that the bilinear transformation preserves the cross ratio of four points.

OR

9. a) Find the bilinear transformation which maps $0, i, \infty$ onto $1, -i, -1$.
 b) Show that $W = \frac{2z + 3}{z - 4}$ maps the circle $x^2 + y^2 - 4x = 0$ onto the straight line $4u + 3 = 0$.



PART – C

Answer **two full** questions.

(2×10=20)

- 10 a) Find the real root of $xe^x - 2 = 0$ by using Regula – Falsi method correct to three decimal places in (0,1).
b) Using Newton-Raphson's method, find the cube root of 37.

OR

- 11 a) Solve $10x + y + z = 12$; $2x + 10y + z = 13$; $2x + 2y + 10z = 14$ by Jacobi iteration method.

- b) By using power method, find the largest eigenvalue of the matrix

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{bmatrix} \text{ given } x_0 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \text{ in five steps.}$$

- 12 a) Using Taylor's series method, find the solution of $\frac{dy}{dx} = x - y^2$, $y(0) = 1$ at $x = 0.2$.

- b) Solve $\frac{dy}{dx} = x + y$ with $x_0 = 0$, $y_0 = 1$ for $x = 0(0.05) 0.05$ using Euler's modified method.

OR

- 13 a) Find the approximate solution at $x = 1.2$ of the equation $\frac{dy}{dx} = xy$, given $y(1) = 2$ by Runge Kutta method by taking $h = 0.2$.

- b) Solve $\frac{dy}{dx} = 1 + \frac{y}{x}$ with $y(1) = 2$, find $y(1.4)$ taking $h = 0.4$ by Euler's modified method.
-